Multiphoton microwave ionization of Li Rydberg atoms

Josh Gurian
Professor Thomas Gallagher

April 28, 2008

Outline

Rydberg atom overview

- $W = \frac{-1}{2n^2}$
- $r \propto n^2$
- $Lifetime \propto n^3$
- $\omega_{kepler} \propto 1/n^3$
- $V_{coulomb} = -\frac{1}{|z|}$



Rydberg atom overview

- $W = \frac{-1}{2n^2}$
- $r \propto n^2$
- $Lifetime \propto n^3$
- $\omega_{kepler} \propto 1/n^3$
- $V = -\frac{1}{|z|} + Ez$



Field Ionization



- $V = -\frac{1}{|z|} + Ez$
- $\frac{dV}{dz} = 0$

Field Ionization



- $V = -\frac{1}{|z|} + Ez$
- $\frac{dV}{dz} = 0$
- $V = -2\sqrt{E}$

Field Ionization



- $V = -\frac{1}{|z|} + Ez$
- $\frac{dV}{dz} = 0$
- $V = -2\sqrt{E}$
- $E = \frac{W^2}{4}$

Field Ionization



- $E = \frac{W^2}{4}$
- $W = \frac{-1}{2n^2} + \frac{3}{2}nkE$

Field Ionization



- $E = \frac{W^2}{4}$
- $W = \frac{-1}{2n^2} + \frac{3}{2}nkE$
- $E = \frac{1}{9n^4}$

Scaled Frequency

Important characteristic: $\omega_0 = \dfrac{\omega}{\omega_{kepler}} = \omega n^3$

Previous work for $\omega_0 \leq 1$:

- van Leeuwen et al., PRL 55 (1985)
- Jensen et al., PRL 62 (1989)
- Pillet et al., PRA 30 (1984)
- Noel et al., PRA 62 (2000)

Microwave Ionization



Pillet et al.,
PRA 30 (1984)

Photoionization limit?

What happens as we approach the photoionization limit, when $\omega_0 \longrightarrow n$?

Cartoon


MW Ionization Cartoon
Normalized electron signal (arb)
-300
-250
-200
-150
-100
-50
0
Binding energy (GHz)

Cartoon


MW Ionization Cartoon
Normalized electron signal (arb)
-107.1
-89.25
-71.4
-53.55
-35.7
-17.85
0
17.85
Binding energy (GHz)

Cartoon


MW Ionization Cartoon
Normalized electron signal (arb)
No MW
0.11 mV/cm
-107.1
-89.25
-71.4
-53.55
-35.7
-17.85
0
17.85
Binding energy (GHz)

Cartoon


MW Ionization Cartoon
Normalized electron signal (arb)
No MW
0.11 mV/cm
0.67 mV/cm
-107.1
-89.25
-71.4
-53.55
-35.7
-17.85
0
17.85
Binding energy (GHz)

Cartoon


MW Ionization Cartoon
Normalized electron signal (arb)
No MW
0.11 mV/cm
0.67 mV/cm
1.83 mV/cm
-107.1
-89.25
-71.4
-53.55
-35.7
-17.85
0
17.85
Binding energy (GHz)

Cartoon


MW Ionization Cartoon
Normalized electron signal (arb)
No MW
0.11 mV/cm
0.67 mV/cm
1.83 mV/cm
3.8 mV/cm
-107.1
-89.25
-71.4
-53.55
-35.7
-17.85
0
17.85
Binding energy (GHz)

Cartoon


MW Ionization Cartoon
Normalized electron signal (arb)
No MW
0.11 mV/cm
0.67 mV/cm
1.83 mV/cm
3.8 mV/cm
6.6 mV/cm
-107.1
-89.25
-71.4
-53.55
-35.7
-17.85
0
17.85
Binding energy (GHz)

Outline

Coherent Evolution-30 Nd:YLF Laser



Coherent Evolution-30 Nd:YLF Laser

- 1st Pulse: 20ns
- 2nd Pulse: 35ns
- 3rd Pulse: 41ns



Laser Setup


Amp
Dye Laser
Dye Laser
Dump
3
BS
Pump Laser
PBS
2
BS
PC
PC
1
Dye Laser
PBS

Experimental Setup





Experimental Setup



Timing


Laser
Pulses
100-500 ns MW Pulse
Field Pulse
0-2V/cm
0
200
400
600
800
1000
time (ns)

Experimental Setup





Outline

Cartoon


MW Ionization Cartoon
Normalized electron signal (arb)
No MW
0.11 mV/cm
0.67 mV/cm
1.83 mV/cm
3.8 mV/cm
6.6 mV/cm
-107.1
-89.25
-71.4
-53.55
-35.7
-17.85
0
17.85
Binding energy (GHz)

Microwave Ionization Steps - Powers



Pulsewidths



Ionization Rate



Theoretical Predictions

$$\Gamma_{exp}(n \approx 360, E \approx 0.8 V/cm, 50 ns) = 5.6 \times 10^6 s^{-1}$$

Hoogenraad and Noordam, PRA 57 (1998):

$$< n|r|n' > \approx \frac{0.4108 n^{-3/2} n'^{-3/2}}{\omega^{5/3}}$$

We can calculate the ionization rate: $\Gamma = 2\pi| < n|r|\epsilon > E|^2$

$$\Gamma(n = 360, E = 0.8 V/cm) = 8.2 \times 10^7 s^{-1}$$

$$\boxed{\Gamma(n = 360, E = 0.2 V/cm) = 5.6 \times 10^6 s^{-1}}$$

Bias Field


Bias voltages on top plate for 200ns MW pulse at 17.850 GHz, 1 V/cm
Normalized electron signal (arb)
0 V/cm
8.5 mV/cm
17 mV/cm
85 mV/cm
-178.5
-160.65
-142.8
-124.95
-107.1
-89.25
-71.4
-53.55
-35.7
-17.85
0
Binding energy (GHz)

Final State Distribution

$$E = 1/9n^4$$



Final State Distribution



Final State Distribution

$$E = 1/9n^4$$



Final State Distribution



Summary

- Introduction to Rydberg field ionization
- Overview of microwave ionization
- Experiment setup
- First results for microwave multiphoton ionization of Rydberg Li atoms

Thanks

Professor Tom Gallagher
Dr. Haruka Maeda
Dr. Ed Shuman
Paul Tanner
Jianing Han
Jirakan Nunkaew
Don Norum
Hyunwook Park
National Science Foundation

Final State Distribution Comparison





Dye laser output



Atomic Conductance

Schelle, Delande, and Buchleitner: